Not an actual patient.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

For adults with idiopathic macular telangiectasia type 2, also known as MacTel

Getting started with ENCELTO

Understanding the process from prescription to procedure

IMPORTANT SAFETY INFORMATION

Who should not receive an ENCELTO surgical implant?

ENCELTO has not been tested in pediatric patients or pregnant women.

The outpatient surgical procedure should not be performed if you are currently experiencing an active or suspected eye infection.

You should not receive ENCELTO if you have a known hypersensitivity to endothelial serum-free media (Endo-SFM).

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

Before receiving ENCELTO, tell your eye surgeon about all your medical conditions, including:

- Are pregnant or plan to become pregnant. Although studies have shown that rhCNTF does not enter the bloodstream, its effects on an unborn baby have not been fully studied.
- Are breastfeeding or plan to breastfeed. It is not known if rhCNTF passes into your breast milk.
- Any current infections.
- Are currently taking or have recently taken medicines that lower the chance of blood clots forming in the body, such as warfarin, low or regular doses of aspirin, or nonsteroidal anti-inflammatory drugs (NSAIDs).

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

About this brochure

You've made the decision to move forward with ENCELTO treatment. This brochure will give you a roadmap through the process, with information on 3 key topics, to help you understand what to expect.

Insurance insights & financial options

How ENCELTOconnect can help 4-5

Information about cost and coverage 6-9

Scheduling your procedure

How to schedule your procedure 10-11

Surgery & post-procedure care

What to expect on procedure day 12-13

Guidance for post-procedure care 14-15

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

ENCELTO connect™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

A support team is ready to help you

ENCELTOconnect is a support program for people who have been prescribed ENCELTO.

- To enroll, **your retina specialist's office will help you complete an enrollment form**. You must sign the form. If the form is not signed when it is first submitted, ENCELTOconnect will contact you to complete an electronic signature

Once you have enrolled, **you'll receive personalized, 1-on-1 help from a Patient Access Consultant (PAC)**.

- Your PAC will reach out to you within 24 to 48 hours of receiving the completed enrollment form from your retina specialist's office. Most of the time, you'll speak with the same PAC, but you might hear from someone else at times if your PAC is unavailable

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

Not an actual patient.

You may not recognize the PAC's phone number, so be sure to answer any unknown calls during this time.

My PAC's name:

_____

Notes:

The next few pages provide an overview of specific ways your PAC can help you prepare for your ENCELTO procedure.

If you have any questions or want to talk to your PAC, you can call **ENCELTOconnect at 1-877-ENCELTO (1-877-362-3586)** Mon-Fri, 8 AM-8 PM EST, and ask for them by name. It's a good idea to save this ENCELTOconnect number in your phone.

IMPORTANT SAFETY INFORMATION (cont'd)

How is ENCELTO administered?

ENCELTO is inserted into the eye as an outpatient surgical procedure performed by an eye surgeon experienced in vitreoretinal surgery. If removal of ENCELTO is necessary, the removal surgery must also be done by an eye surgeon experienced in vitreoretinal surgery in an operating room as an outpatient surgery.

ENCELTO connect™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

Finding out about coverage and costs

Your PAC will provide support based on your unique needs, such as:

Insurance insights

Check coverage and provide guidance on your benefits

Financial assistance options

Explore eligibility for co-pay and travel assistance

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

Not an actual patient.

Coverage insights provided in 24 to 48 hours

The ENCELTO procedure can only be scheduled after the insurance approval process is done. This can take time. Your PAC will reach out on a regular basis to keep you updated on where you stand in the approval process. If you have any questions, you are always welcome to call your PAC.

IMPORTANT SAFETY INFORMATION (cont'd)

What should I avoid after placement of ENCELTO?

Immediately postoperative:

- Avoid heavy lifting (over 20 pounds) for 1 week.
- Keep water out of the eye (eg, close your eye while showering) for 1 week.
- Protect your eyes by wearing glasses or protective eyewear during the day and using an eye shield at night for 1 week.
- Do not drive or use machinery until the eye shield has been removed and your eye surgeon informs you that your vision has recovered to an acceptable level.

ENCELTO connect™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

$0 co-pay assistance*

ENCELTOconnect offers a co-pay assistance program for eligible, commercially insured patients. This program can help with co-pays, out-of-pocket deductibles, or co-insurance expenses related to ENCELTO. (Healthcare provider and procedure-related costs are not included.) If you are eligible, you will be automatically enrolled in the program.

After you join ENCELTOconnect, your PAC will contact your insurance company to find out how both ENCELTO and the surgical procedure are covered. Your PAC will let you know what costs for ENCELTO you might need to pay and if you qualify for co-pay assistance. Eligible patients with commercial insurance may pay $0 for ENCELTO.

The surgical center will let you know about any costs for the procedure itself.

If you receive a letter stating you're eligible for co-pay assistance, bring the letter to your appointment on the day of the procedure.

*For eligible, commercially insured patients. Healthcare provider and procedure-related costs not included.

IMPORTANT SAFETY INFORMATION (cont'd)

What should I avoid after placement of ENCELTO? (cont'd)

Postoperative care:

- Use a topical antibiotic solution at a frequency of 1 drop 4 times a day for 7 days.
- Use a steroid drop taper of prednisolone acetate 1% (or equivalent) starting the day after surgery with the following taper:
  - 1 drop 4 times a day for the first 7 days;
  - 1 drop 3 times a day for the next 7 days;
  - 1 drop 2 times a day for the next 7 days;
  - 1 drop once a day for the last 7 days.

Travel-related assistance

ENCELTOconnect also offers travel-related financial assistance to eligible patients who need to travel a long distance for their surgery. Travel assistance is subject to terms and conditions.

Insurance assistance

It is not unusual for the insurance company to first request more information from your retina specialist's office before providing coverage. Your PAC will provide education to the office to appeal any insurance denials.

We are committed to helping you get access to treatment. Your PAC will help you understand your insurance coverage and explore assistance options.

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

Scheduling your procedure

After the insurance coverage approval process is completed, your retina specialist's office will reach out to schedule your procedure. You can write down all the information on the next page as a handy reminder.

The steps of the procedure may vary because retina specialists can approach the surgery in different ways. Ask your retina specialist what you can expect and write the answers in the "Notes" section to the right.

You must have someone take you home after the procedure. Make sure to include your friend, family member, or caregiver when scheduling your procedure.

IMPORTANT SAFETY INFORMATION (cont'd)

Magnetic Resonance (MR) Conditional Information

IMPORTANT: ENCELTO is MR Conditional

You will receive an implant card, which should be shown to your imaging technician if you need magnetic resonance imaging (MRI) at any time while the ENCELTO implant is in your eye. The card will include details about the ENCELTO implant, the date of insertion, and on the back, instructions for the imaging technician to access important MRI safety information.

What are the possible side effects of ENCELTO?

Please follow all postoperative instructions given by your eye surgeon and ensure you attend all follow-up visits as recommended.

Details about my ENCELTO procedure

Date: _____

Time: _____

Location: _____

My driver: _____

Notes:

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

What you can expect

Day of the procedure:

- Surgery will take place at your designated outpatient surgical center—not your retina specialist's office
- Bring the letter about co-pay assistance (if you have one) to your appointment
- Your retina specialist will tell you how long the procedure may take. In some cases, it may take as little as 45 minutes, after which, you can go home
- You won't be able to leave without assistance, so be sure to have someone ready to take you home after the procedure

IMPORTANT SAFETY INFORMATION (cont'd)

Potential side effects:
Please be advised that ENCELTO and the surgical insertion have related risks such as, but not limited to, endophthalmitis (eye infection), retinal tear and detachment (retina tears and potentially separates from the eye wall resulting in vision loss), vitreous hemorrhage (bleeding within the central cavity of the eye), implant extrusion (the ENCELTO begins to work its way out of the eye), suture-related issues (such as suture-related eye irritation or exposure of sutures), temporary or permanent loss of vision, accelerated cataract formation (clouding of the lens of the eye), and delayed dark adaptation (the ability of the eye to adjust from bright lighting conditions to dark lighting conditions).

Not an actual patient.

After the procedure day:

- Your retina specialist will provide aftercare instructions and help schedule follow-up appointments
  - Follow all aftercare instructions (please see page 14 for more information)
  - Go to your follow-up appointments

If you have any questions about the procedure, ask your retina specialist.

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

Your aftercare instructions

Your retina specialist will talk with you about the recovery process after the ENCELTO procedure and provide specific guidance to follow. Keep reading for some general information that you may find helpful.

Immediately post-procedure:

- Avoid heavy lifting (over 20 pounds) for 1 week
- Keep water out of the eye (eg, close your eye while showering) for 1 week
- Protect your eyes by wearing glasses or protective eyewear during the day and using an eye shield at night for 1 week
- Do not drive or use machinery until the eye shield has been removed and your retina specialist informs you that your vision has recovered to an acceptable level

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

Post-procedure care:

- Use a **topical antibiotic solution** at a frequency of 1 drop, 4 times a day, for 7 days
- Use a **steroid drop** starting the day after surgery with the following taper:

First 7 days: 1 drop, 4 times a day

Next 7 days: 1 drop, 3 times a day

Next 7 days: 1 drop, 2 times a day

Last 7 days: 1 drop, once a day

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

What is ENCELTO?

ENCELTO is an encapsulated cell-based gene therapy. It is a small capsule, about the size of a grain of rice, that is placed inside the eye to release a protein called recombinant human ciliary neurotrophic factor (rhCNTF) that can directly reach the retina, the light-sensitive part of the eye. The capsule contains living cells that have been genetically modified to continuously produce and release rhCNTF. This protein helps protect certain cells in your retina, supporting their health and reducing the loss of light-sensing cells known as photoreceptors.

ENCELTO is used to treat adults with idiopathic macular telangiectasia type 2 (MacTel), a retinal disease that causes progressive vision loss. Your eye surgeon will assess your vision and review your medical history to determine if ENCELTO is the right treatment for you.

IMPORTANT SAFETY INFORMATION

Who should not receive an ENCELTO surgical implant?

ENCELTO has not been tested in pediatric patients or pregnant women.

The outpatient surgical procedure should not be performed if you are currently experiencing an active or suspected eye infection.

You should not receive ENCELTO if you have a known hypersensitivity to endothelial serum-free media (Endo-SFM).

Before receiving ENCELTO, tell your eye surgeon about all your medical conditions, including:

- Are pregnant or plan to become pregnant. Although studies have shown that rhCNTF does not enter the bloodstream, its effects on an unborn baby have not been fully studied.
- Are breastfeeding or plan to breastfeed. It is not known if rhCNTF passes into your breast milk.
- Any current infections.
- Are currently taking or have recently taken medicines that lower the chance of blood clots forming in the body, such as warfarin, low or regular doses of aspirin, or nonsteroidal anti-inflammatory drugs (NSAIDs).

How is ENCELTO administered?

ENCELTO is inserted into the eye as an outpatient surgical procedure performed by an eye surgeon experienced in vitreoretinal surgery. If removal of ENCELTO is necessary, the removal surgery must also be done by an eye surgeon experienced in vitreoretinal surgery in an operating room as an outpatient surgery.

What should I avoid after placement of ENCELTO?

Immediately postoperative:

- Avoid heavy lifting (over 20 pounds) for 1 week.
- Keep water out of the eye (eg, close your eye while showering) for 1 week.
- Protect your eyes by wearing glasses or protective eyewear during the day and using an eye shield at night for 1 week.
- Do not drive or use machinery until the eye shield has been removed and your eye surgeon informs you that your vision has recovered to an acceptable level.

Postoperative care:

- Use a topical antibiotic solution at a frequency of 1 drop 4 times a day for 7 days.
- Use a steroid drop taper of prednisolone acetate 1% (or equivalent) starting the day after surgery with the following taper:
  - 1 drop 4 times a day for the first 7 days;
  - 1 drop 3 times a day for the next 7 days;
  - 1 drop 2 times a day for the next 7 days;
  - 1 drop once a day for the last 7 days.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

Magnetic Resonance (MR) Conditional Information

IMPORTANT: ENCELTO is MR Conditional

You will receive an implant card, which should be shown to your imaging technician if you need magnetic resonance imaging (MRI) at any time while the ENCELTO implant is in your eye. The card will include details about the ENCELTO implant, the date of insertion, and on the back, instructions for the imaging technician to access important MRI safety information.

What are the possible side effects of ENCELTO?

Please follow all postoperative instructions given by your eye surgeon and ensure you attend all follow-up visits as recommended.

Potential side effects:

Please be advised that ENCELTO and the surgical insertion have related risks such as, but not limited to, endophthalmitis (eye infection), retinal tear and detachment (retina tears and potentially separates from the eye wall resulting in vision loss), vitreous hemorrhage (bleeding within the central cavity of the eye), implant extrusion (the ENCELTO begins to work its way out of the eye), suture-related issues (such as suture-related eye irritation or exposure of sutures), temporary or permanent loss of vision, accelerated cataract formation (clouding of the lens of the eye), and delayed dark adaptation (the ability of the eye to adjust from bright lighting conditions to dark lighting conditions).

If delayed dark adaptation occurs, it is unknown for how long these symptoms will be experienced. Take the following safety precautions:

- Driving: Delayed dark adaptation may impair one's ability to see objects, pedestrians, or road signs when moving rapidly from a brightly lit environment to a dimly lit environment (eg, entering a tunnel during the daytime).
- Navigating in the dark: Take caution when moving from bright to dark areas, such as entering a dark room or stepping outside at dusk. Consider using flashlights, nightlights, or motion-activated lighting at home.
- Consider wearing sunglasses or tinted lenses in bright environments to reduce the impact of transitioning from light to dark.

It is common to experience the following symptoms following ENCELTO surgery:

- Sensation of something in the eye (ie, foreign body sensation)
- Eye redness
- Eye irritation
- Eye dryness
- Eye discharge
- Mild to moderate eye pain or discomfort
- Floaters (small spots or shapes that appear in your vision)
- Headache

When to Seek Eye Surgeon Advice

You are to seek immediate care from an eye surgeon if there are sudden changes in your vision, such as an increase in floaters, the appearance of "spider webs," flashing lights, sensitivity to light, loss of vision or visual field, progressively worsening eye pain, or increasing discharge/drainage from the eye as these symptoms could be a sign of a more serious issue.

Call your eye surgeon for medical advice about side effects.
You may report side effects to the Food and Drug Administration (FDA) at 1-800-FDA-1088.

Please see full Prescribing Information.

ENCELTO™
(revakinagene taroretcel-lwey)
implant, for intravitreal use

You're already on the path to slowing MacTel progression

The MacTel journey may feel overwhelming, but you're not alone. Remember, your PAC is ready to help you get started and guide you through insurance coverage, eligibility requirements, and financial assistance options—so you don't have to navigate these things on your own.

Want a quick refresher on ENCELTO before your procedure?

Visit ENCELTO.com for info on what ENCELTO is and how it works.

If you have questions, please call **1-877-ENCELTO (1-877-362-3586)** Mon-Fri, 8 AM-8 PM EST, to speak to a PAC.

IMPORTANT SAFETY INFORMATION (cont'd)

When to Seek Eye Surgeon Advice

You are to seek immediate care from an eye surgeon if there are sudden changes in your vision, such as an increase in floaters, the appearance of "spider webs," flashing lights, sensitivity to light, loss of vision or visual field, progressively worsening eye pain, or increasing discharge/drainage from the eye as these symptoms could be a sign of a more serious issue.

Please see Important Safety Information on pages 16-19 and full Prescribing Information.

neurotech

© Neurotech Pharmaceuticals, Inc. 2026 All rights reserved. US-EO-PM-260400003 05/2026